

CD119 Recombinant Protein

|--|

BackGround:

IFN-γ plays key roles in both the innate and adaptive immune response. IFN- γ activates the cytotoxic activity of innate immune cells, such as macrophages and NK cells. IFN-y production by NK cells and antigen presenting cells (APCs) promotes cell-mediated adaptive immunity by inducing IFN- γ production by T lymphocytes, increasing class I and class II MHC expression, and enhancing peptide antigen presentation. The anti-viral activity of IFN- γ is due to its induction of PKR and other regproteins. Binding of IFN-γ ulatory to the IFNGR1/IFNGR2 complex promotes dimerization of the receptor complexes to form the (IFNGR1/IFNGR2)2 -IFN-y dimer. Binding induces a conformational change in receptor intracellular domains and signaling involves Jak1, Jak2, and Stat1. The critical role of IFN-y in amplification of immune surveillance and function is supported by increased susceptibility to pathogen infection by IFN-y or IFNGR knockout mice and in humans with inactivating mutations in IFNGR1 or IFNGR2. IFN-y also appears to have a role in atherosclerosis.

	11010.
Product:	For researc
PBS, 4M Urea, PH7.4	
Molecular Weight:	
~30kDa	
Swiss-Prot:	

P15260

Purification&Purity:

Transferred into competent cells and the supernatant was purified by NI column affinity chromatography and the purity is > 85% (by SDS-PAGE).

Restriction Sites:

NdeI-XhoI

Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-20 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.

Expression Vector:

pet-22b(+)

DATA:

Da	\sim	S
116		-
6.2	-	
45	-	
35		
25	-	
8.4	-	
4.4	-	and the second

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.				
Add:	1660 South Highway 100, Suite 500 St. Louis Park,			
	MN 55416,USA.			
Email:	info@bioworlde.com			
Tel:	6123263284			
Fax:	6122933841			