

## **CD360 Recombinant Protein**

Catalog:	NCP0308	Host:	E.coli	Tag:	His-tag
BackGround:				Q9HBE5	
The IL-21 receptor (also designated IL-21R, NILR or				Purification&Purity:	
novel interleukin receptor) is a type I cytokine receptor				Transferred into competent cells and the supernatant was	
that forms a complex with the cytokine receptor $\boldsymbol{\gamma}$ chain,				purified by NI column affinity chromatography and the	
$\gamma c$ and mediates IL-21 signaling. IL-21R is present on the				purity is > 85% (by SDS-PAGE).	
surface of natural killer, B and T cell populations with				Restriction Sites:	
high levels in spleen and thymus. IL-21 and IL-21R in-				NdeI-XhoI	
fluence lymphoid proliferation and early lymphoid de-				Storage&Stability:	
velopment in the transition between innate and adaptive immunity. Tumor necrosis factor (TNF) upregulates IL-21R, and combinations of TNF and IL-21 can have synergistic effects on myeloma cell proliferation through pathways involving phosphorylation of JAK1, Stat3 and			and adaptive	Store at 4 ${\rm C}$ short term. Aliquot and store at -20 ${\rm C}$ long	
			upregulates	term. Avoid freeze-thaw cycles.	
			21 can have	Expression Vector: pet-22b(+)	
			ation through		
				DATA:	
16p12.1 an	Erk1/2. The human IL-21R gene maps to chromosome 16p12.1 and encodes a 538 amino acid protein that is closely related to human IL2RB and shares 62% sequence dentity to mouse II21r.			KDa M S 116 66.2	
•				45	
Product: PBS, 4M Urea, PH7.4					
				18.4	
				14.4	
Molecular Weight:				SDS-PAGE Result	
~31kDa				Note:	
Swiss-Prot:				For research use only, not for use in diagnostic procedure.	

## Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: <u>info@biogot.com</u> Tel: 0086-025-68037686 Fax: 0086-025-68035151